

PSYCHOSOCIAL PROBLEMS OF SIGNIFICANT FAMILY MEMBERS OF MENTALLY ILL PATIENTS

Sandhyarani Behera¹, Jitendra Kar², Sagarika Swain³

¹*Department of Psychiatric Nursing, Mental Health Institute (COE), SCBMCH, Cuttack. Mangalabag, Cuttack, Odisha, India.*

^{2,3}*Mental Health Institute (COE), SCBMCH Cuttack. Mangalabag, Cuttack, Odisha, India.*

Corresponding Author -- **Sandhyarani Behera**

Email: sandhyaranibehera2013@yahoo.in

Abstract :

A descriptive study with quantitative approach was undertaken on 50 significant family members of mentally ill patients selected by non-probability convenient sampling technique at Mental Health Institute (COE), SCBMCH, Cuttack to assess the psychosocial problems of significant family members of mentally ill patients. Data was collected through questionnaire on psychosocial problems formulated in the form of 4-point likert scale. Collected data were analyzed by using descriptive and inferential statistics. Findings revealed that Highest Percentage (40%) of the family members were in the age group of 48–60 years. A majority (66%) of them were male and (92%) of them were Hindus and (8%) of them were Muslim. Majority (60%) of them were married (36%) of them were farmer. Highest percentage (30%) of them were illiterate and majority (50%) of them were having income \leq Rs.5000 and (56%) of them from nuclear family. Highest percentage (58%) of them were from rural area and (44%) of them were mother. Majority (38%) of them had >5 years of illness and (76%) of them were having no family history. Most of the significant family members of mentally ill patients (84%) under this study had moderate problem whereas (8%) of them had mild & also (8%) severe problems.

Keywords - Psychosocial problems, significant family members, mentally ill patients

Received: October 22, 2025; Revised: November 29, 2025; Accepted: December 14, 2025

1. Introduction

India is a secular and pluralistic society characterized by tremendous cultural and ethnic diversity. In India the family is the most important institution that has survived through the ages. India, like most other less industrialized, traditional, eastern

societies is a collectivist (a sense of harmony, interdependence and concern for others) society that emphasizes family integrity, family loyalty, and family unity. More specifically, collectivism is reflected in greater readiness to cooperate with family members and extended kin on decisions affecting most aspects of life, including career choice, mate selection, and marriage [1].

When we look at the family as a unit, the following features are common across the globe: it is universal, permanent, nucleus of all social relationships, has an emotional basis, has a formative influence over its members, teaches its members as to what is their social responsibility and the necessity for co-operation and follows a social regulation [2].

Mental illness can be defined as a clinical, significant, behavioural or psychological syndrome that occurs in a person and this is normally associated with impairment in one or more important areas in functioning, or an important loss of freedom [3].

The society views the mental illness as something to be ashamed of and usually the members of the society initially recognize most of the mentally ill patient's behaviour deviated from the social norms [4].

Globally, it is estimated that 450 million people are affected by mental disorders at any one time. These include 121 million people with depression, 24 million with schizophrenia and 37 million with dementia [5]. Mental illness accounted for about 12.3 % of the global burden of disease in 2001 and it is estimated that by 2020 unipolar depressive disorders will be the second most important cause of disability. The burden of caring for mentally ill patients falls on the family members who provide all necessary support [6].

Mental illness may cause a variety of psychosocial problems such as decreased quality of life for the patient's family members, as well as increased social distance for the patient and the family caring for the patient. The family members who care for relatives with mental illness report feeling stigmatized as a result of their association with the mentally ill [7].

Thus, psychiatric professionals often view the family members of a patient as people of support because they can act as informants regarding the patient and they can act as co-therapists at home [8]. The family members need to be in an optimal social and psychological state. It is reported that reduced function of one family member contributes to the burden of other members and this in turn leads to other family members assuming a critical attitude towards the patient [9]. Such criticism can in some cases lead to a relapse of the patient's illness or to the family feeling overwhelmed by the patient's disruptive behaviour [7][10].

In a study investigating the links between stigma, depressive symptoms and coping amongst caregivers, it was found that stigma may erode the morale of family caregivers and result in withdrawal from potential supporters[11].

Studies have shown that in taking care of the mentally ill, the family caregivers have to face the stress and burdens which includes: feelings of insecurity sorrow and worry potential harm, aggression and also stigmatization by relatives and friends. Lastly social isolation fear in the future [12].

In the past 50 years, a shift towards community care and the deinstitutionalization of psychiatric patients has resulted in transferring of responsibility and day-to-day care to family members [13]. In part, this shifting of responsibility has been caused by a deficit in community support services. The profound psychosocial, physical, and financial impact on the family of individuals with severe mental illness is comparable to that of persons with other illnesses such as Alzheimer's disease or cancer.

During the ward round with students and other team members the investigator observed the family members of long term hospitalized mentally ill persons such as separation from family and friends and source of gratification as well as loss of personal health control over the life circumstances as compare to the other families.

The investigators also felt to gain a preliminary understanding of the psychosocial problems faced by the family members in caring for mentally ill patient. Family members need to be educated to be open to reviewing and encouraged to updating their knowledge to face with psychosocial problems towards caring for mentally ill, and how to cope up with the stress and in order to make mentally ill person to lead an optimum level of life as an individual in the society.

2. Objectives :

1. To identify the psychosocial problems among significant family members of mentally ill patients.
2. To correlate the psychosocial problems of family members of mentally ill patients with their selected demographic variables.

3. Materials and methods :

Research design and approach:

A descriptive research design and quantitative approach was used to conduct the study.

Setting of the Study:

The study was conducted at MHI (COE), SCBMCH, Cuttack.

Sample and sampling technique:

50 significant family members of mentally ill patients selected by non-probability convenient sampling technique.

3. Description of the Tools :

The tool has three parts i.e. part “A” and part “B”

Part “A” consists of demographic variables of significant family members of mentally ill patients.

Part “B” consists of the questionnaire related to the psychosocial problems of significant family members of mentally ill patients. The questionnaire in this section was formulated in the form of likert scale, which consists of 30 items on a 4 -point Likert scale format (4=always, 3=frequently, 2= sometimes, 1=never). The questionnaire was divided into six subheadings such as financial problem, disruption of routine family activities, disruption of family leisure, disruption of the family interaction, effect on physical health of the others in family, effect on mental health of others in family.

Validity and Reliability:

Validity refers to the degree to which an instrument measures what it suppose to measure. Content validity concern the degree to which an instrument has appropriate sample of items for the construct being measured and adequately covers the construct domain. The content validity of the tool (psychosocial problems) was established from various experts in the field of psychiatric, clinical psychology, psychiatric nurse specialist, and statistician.

Reliability of the tool was tested by test-retest method where co-efficient correlation was to find out ($r = 0.88$), the tool was found to be more reliable.

Ethical Committee approval: Approval taken from Institutional Ethical Committee, S.C.B Medical College, Cuttack.

Data collection procedure:

Formal permission obtained from the Director-Cum-Medical superintendent of the MHI(COE), S.C.B.M.C.H., Cuttack for data collection. Informed consent obtained from the significant family members of mentally ill patients.

Planned data analysis:

The collected data were organized, tabulated and analysed by using descriptive and inferential statistics.

Distribution of significant family members of mentally ill patients according to their demographic variables reveals that Highest Percentage (40%) of the family members were in the age group of 48–60 years. A majority (66%) of them were male and (92%) of them were Hindus and (8%) of them were Muslim. Majority (60%) of them were married (36%) of them were farmer. Highest percentage (30%) of them were illiterate and majority (50%) of them were having income \leq Rs.5000 and (56%) of them from nuclear family. Highest percentage (58%) of them were from rural area

and (44%) of them were mother. Majority (38%) of them had >5 years of illness and (76%) of them were having no family history of psychiatric illness.

Table 1 - Percentage of Scores revealing psychosocial problems

Sl. No.	Psychosocial problem	Psychosocial problem Score	Number	%
1.	No problem	1 - 30	0	0
2.	Mild problem	31 - 60	4	8
3.	Moderate problem	61- 90	42	84
4.	Severe problem	91 - 120	4	8
Total			50	100

Table: 1- Depicts that that 84% of the significant family members of mentally ill patients having moderate problem whereas 8% of them having mild and also 8% of them also having severe problems. Hence it can be interpreted that highest percentage of them having moderate problems

4. Implication :

Nursing Practice:

The mental health care professionals to actively work with the family members of patients with psychiatric illness to decide suitable psychosocial intervention to address their psychosocial problem associated with mental illness, to improve their quality of life and enhance their coping skills

Regular in-service education programme can be conducted to refresh, up-to-date knowledge and skill on psychosocial problems of family members can help the family members in patients care

Nursing education:

Recommendations for nursing education include that psychiatric nurses should receive training on the strengths of family members in supporting their mentally-ill family members

Family interventions should focus on expanding training to patients and key relatives about wellness recovery, skills training, and task sharing of household and self-care chores.

Nursing Administration :

The Administrator should motivate the staff nurses, to learn how to identify the psychosocial problems of family members as well as patients; it will help in providing quality care.

Appropriate teaching skills in the form of problem solving and communication are needed to promote the coping abilities and lessen the psychosocial problems of the family members

Nursing Research :

Many more innovative studies have been conducted to identify the psychosocial problems and how to manage in day to day life .The research design, findings and the tool can be used as avenues for further research. .

This study will serve as a valuable reference material for future investigators.

5. Recommendations :

A large- scale study can be carried out to generalize the findings.

A comparative study can be conducted with other chronic patients in different settings.

6. Conclusion :

From the findings of the present study, it can be concluded that the Most of the significant family members of mentally ill patients (84%) under this study had moderate problem whereas (8%) of them had mild & also (8%) severe problems.

References

1. Heitzman J, Worden RL. Washington: GPO for the Library of Congress; 1995.India:A Country Study.
2. Bhushan V, Sachdev DR. 26th Edition. Allahabad: Kitabmahal Publishers; 2006. The Family. In : Introduction to Sociology; pp. 291–322.
3. Denise.F. Polit and Cheryl Tatano Beck, (2004) “Nursing Research” Lippincott Willans and Wilkins, Philadelphia, 7th edition, P .No. 587, 661

4. Kaplan HI and Sedock BJ, (2000), “*Comprehensive text book of Psychiatry*”, Lippincott Williams and Wilkins, Philadelphia, 7th edition, P .No.2587-2613.
5. WHO. The World Health Report, Mental Health: New Understanding, New Hope. Geneva: WHO; 2001.
6. Mathers CD, Loncar D. Updated Projection of Global Mortality and Burden of Disease,2002–2030: Data source, Methods and Results. WHO; 2006.
7. Ssebunya J, Kigozi F, Lund C, Kizza D, Okello E. BMC International Health and Stakeholder perceptions of mental health stigma and poverty in. *BMC Int Health Hum Rights*. 2009;9:1–9. doi: 10.1186/1472-698X-9-5. [\[PMC free article\]](#) [\[PubMed\]](#) [\[Cross Ref\]](#)
8. Hasui C, Sakamoto S, Sugiura T, Miyata R, Fujii Y. Burden on Family Members of the Mentally Ill :A Naturalistic Study in Japan. *Compr Psychiatry*. 2002; 43(3): 219–222. 10.1053/COMP.2002.32360. [\[PubMed\]](#)
9. Larson JE, Corrigan P. The Stigma of Families with Mental Illness. *Acad Psychiatry* 2008;32:87–91. 10.1176/appi.ap.32.2.87. [\[PubMed\]](#) [\[Cross Ref\]](#)
10. Bøen H, Dalgard OS et al. The importance of social support in the association between psychological distress and somatic health problems and socio-economic factors among older adults living at home : a cross sectional study. *BMC Geriatr*. 2012; 12(1): 1. 10.1186/1471-2318-12-1. [\[PMC free article\]](#) [\[PubMed\]](#) [\[Cross Ref\]](#)
11. Wintersteen RT, Wintersteen LB, Mupedziswa R. Zimbabwean Families of the Mentally Experiences and Support Needs. *J Soc Dev Afr*. 1995;10(1):89–106.
12. Oldridge M, Huches TCI. Psychological well-being in families with a member Suffering from schizophrenia. An investigation into long-standing problems *British Journal of Psychiatry* 1992; 161:249-51.
13. Worried, Tired and Alone...A Report of Mental Health Carers' Issues in WA, 2003.